arXiv:1607.01006v1 [math.CO] 3 Jul 2016

WATSON-TYPE ${}_3F_2$-SERIES AND SUMMATION FORMULAE INVOLVING GENERALIZED HARMONIC NUMBERS

[A,B]CHUANAN WEI

[A]*Department of Mathematics*
Shanghai Normal University, Shanghai 200234, China
[B]*Department of Information Technology*
Hainan Medical College, Haikou 571199, China

ABSTRACT. In terms of the derivative operator and Watson-type ${}_3F_2$-series identities, three families of summation formulae involving generalized harmonic numbers are established.

1. INTRODUCTION

For a complex variable x, define the shifted factorial to be

$$(x)_0 = 0 \quad \text{and} \quad (x)_n = x(x+1)\cdots(x+n-1) \quad \text{with} \quad n \in \mathbb{N}.$$

Following Andrews, Askey and Roy [3, Chapter 2], define the hypergeometric series by

$${}_{1+r}F_s\left[\begin{matrix} a_0, & a_1, & \cdots, & a_r \\ & b_1, & \cdots, & b_s \end{matrix}\,\Big|\, z\right] = \sum_{k=0}^{\infty} \frac{(a_0)_k (a_1)_k \cdots (a_r)_k}{k!(b_1)_k \cdots (b_s)_k} z^k,$$

where $\{a_i\}_{i\geq 0}$ and $\{b_j\}_{j\geq 1}$ are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Then Watson's ${}_3F_2$-series identity (cf. [3, p. 148]) can be stated as

$${}_3F_2\left[\begin{matrix} a, b, c \\ \frac{1+a+b}{2}, 2c \end{matrix}\,\Big|\, 1\right] = \frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(\frac{1}{2}+c)\Gamma(\frac{1-a-b}{2}+c)}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(\frac{1-a}{2}+c)\Gamma(\frac{1-b}{2}+c)}, \tag{1}$$

where $Re(1-a-b+2c) > 0$ and $\Gamma(x)$ is the well-known gamma function

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt \ \text{ with } \ Re(x) > 0.$$

A terminating form of (1) due to Bailey [1] can be expressed as

$${}_3F_2\left[\begin{matrix} a, b, -n \\ \frac{1+a+b}{2}, -2n \end{matrix}\,\Big|\, 1\right] = \frac{(\frac{1+a}{2})_n (\frac{1+b}{2})_n}{(\frac{1}{2})_n (\frac{1+a+b}{2})_n}. \tag{2}$$

For a complex number x and a positive integer ℓ, define generalized harmonic numbers of ℓ-order to be

$$H_0^{\langle \ell \rangle}(x) = 0 \quad \text{and} \quad H_n^{\langle \ell \rangle}(x) = \sum_{k=1}^{n} \frac{1}{(x+k)^{\ell}} \quad \text{with} \quad n \in \mathbb{N}.$$

2010 Mathematics Subject Classification: Primary 05A10 and Secondary 33C20.
Key words and phrases. Hypergeometric series; Watson's ${}_3F_2$-series identity; Derivative operator; Harmonic numbers.
Email address: weichuanan78@ 163.com.

When $x=0$, they become harmonic numbers of ℓ-order

$$H_0^{\langle \ell \rangle}=0 \quad \text{and} \quad H_n^{\langle \ell \rangle}=\sum_{k=1}^{n}\frac{1}{k^{\ell}} \quad \text{with} \quad n\in\mathbb{N}.$$

Fixing $\ell=1$ in $H_0^{\langle \ell \rangle}(x)$ and $H_n^{\langle \ell \rangle}(x)$, we obtain generalized harmonic numbers

$$H_0(x)=0 \quad \text{and} \quad H_n(x)=\sum_{k=1}^{n}\frac{1}{x+k} \quad \text{with} \quad n\in\mathbb{N}.$$

When $x=0$, they reduce to classical harmonic numbers

$$H_0=0 \quad \text{and} \quad H_n=\sum_{k=1}^{n}\frac{1}{k} \quad \text{with} \quad n\in\mathbb{N}.$$

For a differentiable function $f(x)$, define the derivative operator $\mathcal{D}_x$ by

$$\mathcal{D}_x f(x)=\frac{d}{dx}f(x).$$

In order to explain the relation of the derivative operator and generalized harmonic numbers, we introduce the following lemma.

Lemma 1. *Let x and $\{a_j,b_j,c_j,d_j\}_{j=1}^{s}$ be all complex numbers. Then*

$$\mathcal{D}_x\prod_{j=1}^{s}\frac{a_jx+b_j}{c_jx+d_j}=\prod_{j=1}^{s}\frac{a_jx+b_j}{c_jx+d_j}\sum_{j=1}^{s}\frac{a_jd_j-b_jc_j}{(a_jx+b_j)(c_jx+d_j)}.$$

Proof. It is not difficult to verify the case $s=1$ of Lemma 1. Suppose that

$$\mathcal{D}_x\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}=\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}\sum_{j=1}^{m}\frac{a_jd_j-b_jc_j}{(a_jx+b_j)(c_jx+d_j)}$$

is true. We can proceed as follows:

$$\begin{aligned}
\mathcal{D}_x\prod_{j=1}^{m+1}\frac{a_jx+b_j}{c_jx+d_j}&=\mathcal{D}_x\left\{\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}\frac{a_{m+1}x+b_{m+1}}{c_{m+1}x+d_{m+1}}\right\}\\
&=\frac{a_{m+1}x+b_{m+1}}{c_{m+1}x+d_{m+1}}\mathcal{D}_x\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}+\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}\mathcal{D}_x\frac{a_{m+1}x+b_{m+1}}{c_{m+1}x+d_{m+1}}\\
&=\frac{a_{m+1}x+b_{m+1}}{c_{m+1}x+d_{m+1}}\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}\sum_{j=1}^{m}\frac{a_jd_j-b_jc_j}{(a_jx+b_j)(c_jx+d_j)}\\
&+\prod_{j=1}^{m}\frac{a_jx+b_j}{c_jx+d_j}\frac{a_{m+1}d_{m+1}-b_{m+1}c_{m+1}}{(c_{m+1}x+d_{m+1})^2}\\
&=\prod_{j=1}^{m+1}\frac{a_jx+b_j}{c_jx+d_j}\left\{\sum_{j=1}^{m}\frac{a_jd_j-b_jc_j}{(a_jx+b_j)(c_jx+d_j)}+\frac{a_{m+1}d_{m+1}-b_{m+1}c_{m+1}}{(a_{m+1}x+b_{m+1})(c_{m+1}x+d_{m+1})}\right\}\\
&=\prod_{j=1}^{m+1}\frac{a_jx+b_j}{c_jx+d_j}\sum_{j=1}^{m+1}\frac{a_jd_j-b_jc_j}{(a_jx+b_j)(c_jx+d_j)}.
\end{aligned}$$

This proves Lemma 1 inductively. □

Setting $a_j=1,b_j=r-j+1,c_j=0,d_j=j$ in Lemma 1, it is easy to find that

$$\mathcal{D}_x\binom{x+r}{s}=\binom{x+r}{s}\big\{H_r(x)-H_{r-s}(x)\big\},$$

where $r, s \in \mathbb{N}_0$ with $s \leq r$. Besides, we have the following relation:

$$\mathcal{D}_x H_n^{\langle \ell \rangle}(x) = -\ell H_n^{\langle \ell+1 \rangle}(x).$$

As pointed out by Richard Askey (cf. [2]), expressing harmonic numbers in accordance with differentiation of binomial coefficients can be traced back to Issac Newton. In 2003, Paule and Schneider [9] computed the family of series:

$$W_n(\alpha) = \sum_{k=0}^{n} \binom{n}{k}^{\alpha} \{1 + \alpha(n-2k)H_k\}$$

with $\alpha = 1, 2, 3, 4, 5$ by combining this way with Zeilberger's algorithm for definite hypergeometric sums. According to the derivative operator and the hypergeometric form of Andrews' q-series transformation, Krattenthaler and Rivoal [6] deduced general Paule-Schneider type identities with α being a positive integer. More results from differentiation of binomial coefficients can be seen in the papers [11, 15, 16, 18, 19]. For different ways and related harmonic number identities, the reader may refer to [4, 7, 8, 10, 12, 14]. It should be mentioned that Sun [13] showed recently some congruence relations concerning harmonic numbers to us.

Inspired by the work just mentioned, we shall explore, by means of the derivative operator and Watson-type ${}_3F_2$-series identities, closed expressions for the following three families of series:

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} \frac{\binom{2x+n+k}{k}}{\binom{x+k}{k}} k^t H_k,$$
$$\sum_{k=0}^{n} \binom{2n-k}{n} \binom{x+k}{k} k^t H_k^{\langle 2 \rangle}(x),$$
$$\sum_{k=0}^{n} \binom{2n-k}{n} \binom{x+k}{k} k^t H_k^2(x),$$

where $t \in \mathbb{N}_0$. For limit of space, our explicit formulae are offered only for $t = 0, 1, 2$. In order to make the reader have a taste, we enumerate, above all, three concise harmonic number identities from the special cases of them as follows:

$$\sum_{k=0}^{n} (-2)^k \binom{n}{k} H_k = \begin{cases} 2H_{2m} - H_m, & n = 2m; \\ H_m - 2H_{2m+1}, & n = 2m+1, \end{cases}$$
$$\sum_{k=0}^{n} \binom{2n-k}{n} H_k^{\langle 2 \rangle} = \frac{\binom{2n+1}{n}}{2} \big\{ H_{2n+1}^{\langle 2 \rangle} - (H_{2n+1} - H_n)^2 \big\},$$
$$\sum_{k=0}^{n} \binom{2n-k}{n} H_k^2 = \frac{\binom{2n+1}{n}}{2} \big\{ 2H_{n+1}^{\langle 2 \rangle} - H_{2n+1}^{\langle 2 \rangle} + 2H_{n+1}^2 - H_n^2 + H_{2n+1}(H_{2n+1} - 4H_{n+1} + 2H_n) \big\}.$$

2. The first family of summation formulae involving

Lemma 2. *Let a, b and c be all complex numbers. Then*

$$ {}_3F_2\left[\begin{matrix}a,b,c\\ \frac{1+a+b}{2},2c-1\end{matrix}\,\Big|\,1\right]=\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{1+a+b}{2})}{\Gamma(\frac{a}{2})\Gamma(\frac{b}{2})\Gamma(c-\frac{a}{2})\Gamma(c-\frac{b}{2})}$$
$$ +\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{1+a+b}{2})}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(c-\frac{1+a}{2})\Gamma(c-\frac{1+b}{2})}, $$

where the convergence condition is $Re(2c-a-b-1)>0$.

Proof. Recall Whipple's ${}_3F_2$-series identity (cf. [3, p. 149]):

$$ {}_3F_2\left[\begin{matrix}a,1-a,b\\ c,1+2b-c\end{matrix}\,\Big|\,1\right]=\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{a+c}{2})\Gamma(\frac{1+a-c}{2}+b)\Gamma(\frac{1-a+c}{2})\Gamma(\frac{2-a-c}{2}+b)} $$

provided that $Re(b)>0$. Perform the replacement $a\to 1+a$ in the last equation to get

$$ {}_3F_2\left[\begin{matrix}1+a,-a,b\\ c,1+2b-c\end{matrix}\,\Big|\,1\right]=\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(\frac{1-a-c}{2}+b)}. $$

The linear combination of the last two equations gives

$$ {}_3F_2\left[\begin{matrix}a,-a,b\\ c,1+2b-c\end{matrix}\,\Big|\,1\right]=\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{a+c}{2})\Gamma(\frac{1+a-c}{2}+b)\Gamma(\frac{1-a+c}{2})\Gamma(\frac{2-a-c}{2}+b)} $$
$$ +\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(\frac{1-a-c}{2}+b)}. \tag{3} $$

In terms of Kummer's transformation formula (cf. [3, p. 143]):

$$ {}_3F_2\left[\begin{matrix}a,b,c\\ d,e\end{matrix}\,\Big|\,1\right]=\frac{\Gamma(d)\Gamma(e)\Gamma(d+e-a-b-c)}{\Gamma(a)\Gamma(d+e-a-b)\Gamma(d+e-a-c)} $$
$$ \times\ {}_3F_2\left[\begin{matrix}d-a,e-a,d+e-a-b-c\\ d+e-a-b,d+e-a-c\end{matrix}\,\Big|\,1\right], \tag{4} $$

we gain

$$ 3F_2\left[\begin{matrix}c,b,a\\ \frac{1+a+b}{2},2c-1\end{matrix}\,\Big|\,1\right]=\frac{\Gamma(\frac{1+a+b}{2})\Gamma(2c-1)\Gamma(c-\frac{1+a+b}{2})}{\Gamma(c)\Gamma(c-\frac{1+b-a}{2})\Gamma(c-\frac{1+a-b}{2})} $$
$$ \times\ {}_3F_2\left[\begin{matrix}\frac{1+a+b}{2}-c,c-1,c-\frac{1+a+b}{2}\\ c-\frac{1+b-a}{2},c-\frac{1+a-b}{2}\end{matrix}\,\Big|\,1\right]. $$

Calculating the series on the right hand side by (3), we achieve Lemma 2 to complete the proof. □

Theorem 3. *Let x be a complex number. Then*

$$ \sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}}{\binom{x+k}{k}}H_k=\begin{cases}H_m(x)+2H_{2m}-H_m, & n=2m;\\ H_m-2H_{2m+1}-H_m(x), & n=2m+1.\end{cases} $$

Proof. The case $a=-n$, $b=2x+n+1$ and $c=y+1$ of Lemma 2 reads as

$$ \sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}=\begin{cases}\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}, & n=2m;\\ \frac{-1}{2y+1}\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}, & n=2m+1.\end{cases} $$

Applying the derivative operator $\mathcal{D}_y$ to both sides of it, we attain

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}\big\{2H_k(2y)-H_k(y)\big\}$$
$$=\begin{cases}\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}\big\{H_m(x-y)+H_m(y-\frac{1}{2})\big\}, & n=2m;\\ \frac{-1}{2y+1}\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}\big\{H_m(x-y)+H_{m+1}(y-\frac{1}{2})\big\}, & n=2m+1.\end{cases}$$

Choosing $y=0$ in the last equation, we obtain Theorem 3 to finish the proof. □

Taking $x=p$ with $p\in\mathbb{N}_0$ in Theorem 3, we have the summation formula on harmonic numbers.

Corollary 4. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\binom{2p+n+k}{p+k}H_k$$
$$=\binom{2p+n}{p}\begin{cases}2H_{2m}-H_m+H_{p+m}-H_p, & n=2m;\\ H_p-H_{p+m}+H_m-2H_{2m+1}, & n=2m+1.\end{cases}$$

When $x\to\infty$, Theorem 3 reduces to the concise harmonic number identity:

$$\sum_{k=0}^{n}(-2)^k\binom{n}{k}H_k=\begin{cases}2H_{2m}-H_m, & n=2m;\\ H_m-2H_{2m+1}, & n=2m+1.\end{cases}$$

Lemma 5. *Let a, b and c be all complex numbers. Then*

$$\sum_{k=0}^{\infty}k\frac{(a)_k(b)_k(c)_k}{k!(\frac{1+a+b}{2})_k(2c-1)_k}=ab\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{3+a+b}{2})}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(c-\frac{1+a}{2})\Gamma(c-\frac{1+b}{2})}$$
$$+\{2c^2-(a+b+3)c+(a+1)(b+1)\}$$
$$\times\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{3+a+b}{2})}{\Gamma(\frac{a}{2})\Gamma(\frac{b}{2})\Gamma(c-\frac{a}{2})\Gamma(c-\frac{b}{2})},$$

where the convergence condition is $Re(2c-a-b-3)>0$.

Proof. Employ the substitution $a\to 1+a$ in (3) to get

$${}_3F_2\left[\begin{matrix}1+a,-1-a,b\\ c,1+2b-c\end{matrix}\,\middle|\,1\right]=\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(b-\frac{a+c-1}{2})}$$
$$+\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{2+a+c}{2})\Gamma(\frac{3+a-c}{2}+b)\Gamma(\frac{c-a-1}{2})\Gamma(b-\frac{a+c}{2})}.$$

The linear combination of (3) and the last equation provides

$${}_3F_2\left[\begin{matrix}a,-1-a,b\\ c,1+2b-c\end{matrix}\,\middle|\,1\right]=\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(\frac{1-a-c}{2}+b)}$$
$$+\frac{a^2+a+(1+2b-c)c}{(1+a+2b-c)(a+c)}$$
$$\times\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{a+c}{2})\Gamma(\frac{1+a-c}{2}+b)\Gamma(\frac{1-a+c}{2})\Gamma(\frac{2-a-c}{2}+b)}. \tag{5}$$

In accordance with (4), we gain

$$
\begin{aligned}
{}_3F_2\left[\begin{matrix}c,b,a\\ \frac{1+a+b}{2},2c-2\end{matrix}\,\Big|\,1\right] &= \frac{\Gamma(\frac{1+a+b}{2})\Gamma(2c-2)\Gamma(c-\frac{3+a+b}{2})}{\Gamma(c)\Gamma(c-\frac{3+b-a}{2})\Gamma(c-\frac{3+a-b}{2})}\\
&\quad\times {}_3F_2\left[\begin{matrix}\frac{1+a+b}{2}-c,c-2,c-\frac{3+a+b}{2}\\ c-\frac{3+b-a}{2},c-\frac{3+a-b}{2}\end{matrix}\,\Big|\,1\right].
\end{aligned}
$$

Computing the series on the right hand side by (5), we achieve

$$
\begin{aligned}
{}_3F_2\left[\begin{matrix}a,b,c\\ \frac{1+a+b}{2},2c-2\end{matrix}\,\Big|\,1\right] &= \frac{2}{c-1}\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{3+a+b}{2})}{\Gamma(\frac{a}{2})\Gamma(\frac{b}{2})\Gamma(c-\frac{2+a}{2})\Gamma(c-\frac{2+b}{2})}\\
&\quad+\frac{2c^2-(a+b+5)c+ab+a+b+3}{2(c-1)}\\
&\quad\times\frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{3+a+b}{2})}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(c-\frac{1+a}{2})\Gamma(c-\frac{1+b}{2})}. \qquad (6)
\end{aligned}
$$

It is routine to show that

$$
\begin{aligned}
\sum_{k=0}^{\infty}k\frac{(a)_k(b)_k(c)_k}{k!(\frac{1+a+b}{2})_k(2c-1)_k} &= \sum_{k=1}^{\infty}\frac{(a)_k(b)_k(c)_k}{(k-1)!(\frac{1+a+b}{2})_k(2c-1)_k}\\
&= \sum_{k=0}^{\infty}\frac{(a)_{k+1}(b)_{k+1}(c)_{k+1}}{k!(\frac{1+a+b}{2})_{k+1}(2c-1)_{k+1}}\\
&= \frac{2abc}{(1+a+b)(2c-1)}{}_3F_2\left[\begin{matrix}1+a,1+b,1+c\\ \frac{3+a+b}{2},2c\end{matrix}\,\Big|\,1\right].
\end{aligned}
$$

Evaluating the series on the right hand side by (6), we attain Lemma 5 to complete the proof. □

Theorem 6. *Let x be a complex number. Then*

$$
\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}}{\binom{x+k}{k}}kH_k = \frac{n(2x+n+1)}{x+1}\\
&\times\begin{cases}H_{m-1}(x+1)+2H_{2m}-H_m, & n=2m;\\ H_m-2H_{2m}-H_m(x+1)-\frac{x+1}{(2m+1)(x+m+1)}, & n=2m+1.\end{cases}
\end{aligned}
$$

Proof. The case $a=-n$, $b=2x+n+1$ and $c=y+1$ of Lemma 5 can be written as

$$
\begin{aligned}
&\sum_{k=0}^{n}(-1)^k k\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}\\
&=\begin{cases}\frac{2m(2x+2m+1)}{x-y+1}\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}, & n=2m;\\ \frac{2\{y^2-xy-(2m+1)(x+m+1)\}}{(x-y+1)(2y+1)}\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}, & n=2m+1.\end{cases}
\end{aligned}
$$

Applying the derivative operator $\mathcal{D}_y$ to both sides of it, we obtain

$$
\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}k\big\{2H_k(2y)-H_k(y)\big\}\\
&=\begin{cases}\frac{2m(2x+2m+1)}{x-y+1}\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}\big\{H_{m-1}(x-y+1)+H_m(y-\frac{1}{2})\big\}, & n=2m;\\ \frac{2\alpha_m(x,y)}{(x-y+1)(2y+1)}\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}\big\{H_m(x-y)+H_m(y+\frac{1}{2})-\beta_m(x,y)\big\}, & n=2m+1,\end{cases}
\end{aligned}
$$

where the corresponding expressions are

$$\alpha_m(x,y) = y^2 - xy - (2m+1)(x+m+1),$$
$$\beta_m(x,y) = \frac{m(2x+2m+3)(2x-4y+1)+(x-y+1)^2}{(x-y+1)(2y+1)\{y^2-xy-(2m+1)(x+m+1)\}}.$$

Selecting $y=0$ in the last equation, we get Theorem 6 to finish the proof. □

Fixing $x=p$ with $p\in\mathbb{N}_0$ in Theorem 6, we deduce the summation formula on harmonic numbers.

Corollary 7. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\binom{2p+n+k}{p+k}kH_k = n\binom{2p+n+1}{p+1}$$
$$\times\begin{cases}2H_{2m}-H_m+H_{p+m}-H_{p+1}, & n=2m;\\ H_m-2H_{2m}+H_{p+1}-H_{p+m+1}-\frac{p+1}{(2m+1)(p+m+1)}, & n=2m+1.\end{cases}$$

When $x\to\infty$, Theorem 6 produces the concise harmonic number identity:

$$\sum_{k=0}^{n}(-2)^k\binom{n}{k}kH_k = \begin{cases}4m(2H_{2m}-H_m), & n=2m;\\ 2(2m+1)(H_m-2H_{2m+1})+2, & n=2m+1.\end{cases}$$

Lemma 8. *Let a, b and c be all complex numbers. Then*

$$\sum_{k=0}^{\infty}k^2\frac{(a)_k(b)_k(c)_k}{k!(\frac{1+a+b}{2})_k(2c-1)_k} = \Phi(a,b,c)\frac{\Gamma(\frac{3}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{5+a+b}{2})}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(c-\frac{1+a}{2})\Gamma(c-\frac{1+b}{2})}$$
$$+\Psi(a,b,c)\frac{\Gamma(\frac{3}{2})\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac{1}{2})\Gamma(c-\frac{5+a+b}{2})}{\Gamma(\frac{a}{2})\Gamma(\frac{b}{2})\Gamma(c-\frac{a}{2})\Gamma(c-\frac{b}{2})},$$

where the symbols on the right hand side stand for

$$\Phi(a,b,c) = ab\{2c^2-(1+a+b)c+2(a+b+ab-1)\},$$
$$\Psi(a,b,c) = 4c^3+2(a+b+3ab-5)c^2$$
$$-(3a^2b+3ab^2+11ab+2a^2+2a+2b^2+2b-8)c$$
$$+2(1+a)(1+b)(a+b+ab-1)$$

and the parameters satisfy the condition $Re(2c-a-b-5)>0$.

Proof. Replace a by $1+a$ in (5) to gain

$${}_3F_2\left[\begin{matrix}1+a,-2-a,b\\ c,1+2b-c\end{matrix}\,\middle|\,1\right] = \frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{2+a+c}{2})\Gamma(\frac{3+a-c}{2}+b)\Gamma(\frac{c-a-1}{2})\Gamma(b-\frac{a+c}{2})}$$
$$+\frac{a^2+3a+2+(1+2b-c)c}{(2+a+2b-c)(1+a+c)}$$
$$\times\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(\frac{1-a-c}{2}+b)}.$$

The linear combination of (5) and the last equation offers

$$\begin{aligned}
&{}_3F_2\left[\begin{matrix}a,-2-a,b\\ c,1+2b-c\end{matrix}\,\Big|\,1\right]\\
&=\frac{(2+a)(1+a+b)+(1+2b-c)c}{(2+a+2b-c)(1+a+c)}\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{c-a}{2})\Gamma(\frac{1-a-c}{2}+b)}\\
&+\frac{a(1+a-b)+(1+2b-c)c}{(a-2b+c)(1+a-c)}\frac{\pi\Gamma(c)\Gamma(1+2b-c)}{4^b\Gamma(\frac{2+a+c}{2})\Gamma(\frac{3+a-c}{2}+b)\Gamma(\frac{c-a-1}{2})\Gamma(b-\frac{a+c}{2})}.
\end{aligned}\tag{7}$$

According to (4), we achieve

$$\begin{aligned}
{}_3F_2\left[\begin{matrix}c,b,a\\ \frac{1+a+b}{2},2c-3\end{matrix}\,\Big|\,1\right]&=\frac{\Gamma(\frac{1+a+b}{2})\Gamma(2c-3)\Gamma(c-\frac{5+a+b}{2})}{\Gamma(c)\Gamma(c-\frac{5+b-a}{2})\Gamma(c-\frac{5+a-b}{2})}\\
&\times{}_3F_2\left[\begin{matrix}\frac{1+a+b}{2}-c,c-3,c-\frac{5+a+b}{2}\\ c-\frac{5+b-a}{2},c-\frac{5+a-b}{2}\end{matrix}\,\Big|\,1\right].
\end{aligned}$$

Reckoning the series on the right hand side by (7), we attain

$$\begin{aligned}
&{}_3F_2\left[\begin{matrix}a,b,c\\ \frac{1+a+b}{2},2c-3\end{matrix}\,\Big|\,1\right]\\
&=\left\{1+\tfrac{3ab(c-3)}{(c-1)(2c-a-5)(2c-b-5)}\right\}\frac{\Gamma(\frac12)\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac52)\Gamma(c-\frac{5+a+b}{2})}{\Gamma(\frac{1+a}{2})\Gamma(\frac{1+b}{2})\Gamma(c-\frac{5+a}{2})\Gamma(c-\frac{5+b}{2})}\\
&+\tfrac{2\{6c^2-3c(a+b+7)+2ab+5a+5b+17\}}{(c-1)(2c-a-4)(2c-b-4)}\frac{\Gamma(\frac12)\Gamma(\frac{1+a+b}{2})\Gamma(c-\frac32)\Gamma(c-\frac{5+a+b}{2})}{\Gamma(\frac{a}{2})\Gamma(\frac{b}{2})\Gamma(c-\frac{4+a}{2})\Gamma(c-\frac{4+b}{2})}.
\end{aligned}\tag{8}$$

It is easy to see that

$$\begin{aligned}
\sum_{k=0}^{\infty}k^2\frac{(a)_k(b)_k(c)_k}{k!(\frac{1+a+b}{2})_k(2c-1)_k}&=\sum_{k=1}^{\infty}k\frac{(a)_k(b)_k(c)_k}{(k-1)!(\frac{1+a+b}{2})_k(2c-1)_k}\\
&=\sum_{k=0}^{\infty}(k+1)\frac{(a)_{k+1}(b)_{k+1}(c)_{k+1}}{k!(\frac{1+a+b}{2})_{k+1}(2c-1)_{k+1}}\\
&=\frac{2abc}{(1+a+b)(2c-1)}{}_3F_2\left[\begin{matrix}1+a,1+b,1+c\\ \frac{3+a+b}{2},2c\end{matrix}\,\Big|\,1\right]\\
&+\frac{2abc}{(1+a+b)(2c-1)}\sum_{k=1}^{\infty}\frac{(1+a)_k(1+b)_k(1+c)_k}{(k-1)!(\frac{3+a+b}{2})_k(2c)_k}\\
&=\frac{2abc}{(1+a+b)(2c-1)}{}_3F_2\left[\begin{matrix}1+a,1+b,1+c\\ \frac{3+a+b}{2},2c\end{matrix}\,\Big|\,1\right]\\
&+\frac{2ab(1+a)(1+b)(1+c)}{(1+a+b)(3+a+b)(2c-1)}{}_3F_2\left[\begin{matrix}2+a,2+b,2+c\\ \frac{5+a+b}{2},1+2c\end{matrix}\,\Big|\,1\right].
\end{aligned}$$

Calculating the two series on the right hand side by (6) and (8), we obtain Lemma 8 to complete the proof. □

Theorem 9. *Let x be a complex number. Then*

$$\begin{aligned}
\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}}{\binom{x+k}{k}}k^2H_k&=\frac{n(2x+n+1)(n^2+n+2nx-x)}{(x+1)(x+2)}\\
\times&\begin{cases}2H_{2m}-H_m+H_{m-2}(x+2)+\frac{1-x}{2m(2x+2m+1)-x}, & n=2m;\\ H_m-2H_{2m}-H_{m-1}(x+2)-\frac{x+3}{2m(2x+2m+3)+x+2}, & n=2m+1.\end{cases}
\end{aligned}$$

Proof. The case $a=-n$, $b=2x+n+1$ and $c=y+1$ of Lemma 8 can be manipulated as

$$\sum_{k=0}^{n}(-1)^k k^2\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}=\begin{cases}\gamma_m(x,y)\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}, & n=2m;\\ \eta_m(x,y)\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}, & n=2m+1,\end{cases}$$

where the corresponding expressions are

$$\begin{aligned}\gamma_m(x,y)&=\frac{2m(2x+2m+1)\{4m^2+2(2x+1)m+xy-y^2-x-y\}}{(x-y+1)(x-y+2)},\\ \eta_m(x,y)&=\frac{2\{y^2-(m+1)(2m+1)-x(y+2m+1)\}}{(x-y+1)(2y+1)}\\ &\quad+\frac{2m(2x+2m+3)\{3(y+2)(y-x-2)-2(m-1)(2x+2m+5)\}}{(x-y+1)(x-y+2)(2y+1)}.\end{aligned}$$

Applying the derivative operator $\mathcal{D}_y$ to both sides of it, we get

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+n+k}{k}\binom{y+k}{k}}{\binom{x+k}{k}\binom{2y+k}{k}}k^2\big\{2H_k(2y)-H_k(y)\big\}\\ &=\begin{cases}\mu_m(x,y)\frac{\binom{x-y+m}{m}\binom{m-\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m-\frac{1}{2}}{m}}\big\{H_{m-2(x-y+2)}+H_m(y-\frac{1}{2})-\nu_m(x,y)\big\}, & n=2m;\\ \lambda_m(x,y)\frac{\binom{x-y+m}{m}\binom{m+\frac{1}{2}}{m}}{\binom{x+m}{m}\binom{y+m+\frac{1}{2}}{m}}\big\{H_{m-2(x-y+2)}+H_{m+1}(y-\frac{1}{2})-\omega_m(x,y)\big\}, & n=2m+1,\end{cases}\end{aligned}$$

where the symbols on the right hand side stand for

$$\begin{aligned}\mu_m(x,y)&=\frac{2m(2x+2m+1)\{x(y+4m-1)-y(y+1)+2m(2m+1)\}}{(x-y+1)(x-y+2)},\\ \nu_m(x,y)&=\frac{x-2y-1}{x(y+4m-1)-y(y+1)+2m(2m+1)},\\ \lambda_m(x,y)&=\frac{\begin{cases}2m(2x+2m+3)\{3y^2-3xy-2(2m+1)(x+m+1)\}\\ +2(x-y+2)\{y^2-xy-(2m+1)(x+m+1)\}\end{cases}}{(x-y+1)(x-y+2)(2y+1)},\\ \omega_m(x,y)&=\frac{m(3x-6y-1)(2x+2m+3)+x^2+x-1-4(x+1)y+3y^2}{\begin{cases}(2m+1)(x+m+1)(2+x+4mx+6m+4m^2)\\ -(1+3m+2m^2-x-x^2-6m^2x-7mx-6mx^2)y\\ -(2+9m+6m^2+2x+6mx)y^2+y^3\end{cases}}.\end{aligned}$$

Setting $y=0$ in the last equation, we gain Theorem 9 to finish the proof. □

Choosing $x=p$ with $p\in\mathbb{N}_0$ in Theorem 9, we have the summation formula on harmonic numbers.

Corollary 10. *Let p be a nonnegative integer. Then*

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\binom{2p+n+k}{p+k}k^2H_k=\frac{n(n^2+n+2pn-p)}{p+2}\binom{2p+n+1}{p+1}\\ &\times\begin{cases}2H_{2m}-H_m+H_{p+m}-H_{p+2}+\frac{1-p}{2m(2p+2m+1)-p}, & n=2m;\\ H_m-2H_{2m}+H_{p+2}-H_{p+m+1}-\frac{p+3}{2m(2p+2m+3)+p+2}, & n=2m+1.\end{cases}\end{aligned}$$

When $x\to\infty$, Theorem 9 creates the concise harmonic number identity:

$$\sum_{k=0}^{n}(-2)^k\binom{n}{k}k^2H_k=2n(2n-1)\times\begin{cases}2H_{2m}-H_m-\frac{1}{4m-1},& n=2m;\\ H_m-2H_{2m}-\frac{1}{4m+1},& n=2m+1.\end{cases}$$

3. The second family of summation formulae involving generalized harmonic numbers

Lemma 11. *Let a and b be both complex numbers. Then*

$${}_3F_2\left[\begin{matrix}a,b-a,-n\\ \frac{b}{2},-2n\end{matrix}\,\middle|\,1\right]=\frac{a}{b}\frac{(\frac{2+a}{2})_n(\frac{1+b-a}{2})_n}{(\frac{1}{2})_n(\frac{2+b}{2})_n}+\frac{b-a}{b}\frac{(\frac{1+a}{2})_n(\frac{2+b-a}{2})_n}{(\frac{1}{2})_n(\frac{2+b}{2})_n}.$$

Proof. Perform the replacement $b\to 1+b-a$ in (2) to achieve

$${}_3F_2\left[\begin{matrix}a,1+b-a,-n\\ 1+\frac{b}{2},-2n\end{matrix}\,\middle|\,1\right]=\frac{(\frac{1+a}{2})_n(\frac{2+b-a}{2})_n}{(\frac{1}{2})_n(\frac{2+b}{2})_n}.$$

Replace a by $1+a$ in it to attain

$${}_3F_2\left[\begin{matrix}1+a,b-a,-n\\ 1+\frac{b}{2},-2n\end{matrix}\,\middle|\,1\right]=\frac{(\frac{2+a}{2})_n(\frac{1+b-a}{2})_n}{(\frac{1}{2})_n(\frac{2+b}{2})_n}.$$

The linear combination of the last two equations gives Lemma 11. □

Theorem 12. *Let x be a complex number. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}H_k^{\langle 2\rangle}(x)=\frac{\binom{x+2n+1}{n}}{8}$$
$$\times\left\{4H_{2n}^{\langle 2\rangle}(x+1)-\left[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\right]\left[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})-\tfrac{4}{x+1}\right]\right\}.$$

Proof. The case $a=x+1$ and $b=y+2$ of Lemma 11 reads as

$$\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}=\frac{4^n(x+1)}{y+2}\frac{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}}{\binom{\frac{y+2}{2}+n}{n}}$$
$$+\frac{4^n(y-x+1)}{y+2}\frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}}{\binom{\frac{y+2}{2}+n}{n}}.\tag{9}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (9), we obtain

$$\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}\left\{H_k(x)-H_k(y-x)\right\}=\frac{4^n}{y+2}\frac{1}{\binom{\frac{y+2}{2}+n}{n}}$$
$$\times\left\{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}-\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}+\theta_n(x,y)\right\},$$

where the corresponding expression is

$$\theta_n(x,y)=\frac{(x+1)\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}\left[H_n(\frac{x+1}{2})-H_n(\frac{y-x}{2})\right]}{2}$$
$$+\frac{(y-x+1)\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}\left[H_n(\frac{x}{2})-H_n(\frac{y-x+1}{2})\right]}{2}.$$

The last equation can be reformulated as

$$\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}\sum_{i=1}^{k}\frac{1}{(x+i)(y-x+i)}$$
$$=\frac{4^n}{y+2}\frac{1}{\binom{\frac{y+2}{2}+n}{n}}\left\{\frac{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}-\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}}{y-2x}+\frac{\theta_n(x,y)}{y-2x}\right\}.$$

Finding the limit $y\to 2x$ of it and using the two relations

$$\begin{aligned}
&\mathrm{Lim}_{y\to 2x}\frac{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}-\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}}{y-2x}\\
&=\mathrm{Lim}_{y\to 2x}\mathcal{D}_y\left\{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}-\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}\right\}\\
&=\frac{1}{2}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n}\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big],\\
&\mathrm{Lim}_{y\to 2x}\frac{\theta_n(x,y)}{y-2x}\\
&=\mathrm{Lim}_{y\to 2x}\mathcal{D}_y\theta_n(x,y)\\
&=\frac{1}{2}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n}\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]\\
&+\frac{x+1}{4}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n}\Big\{4H_{2n}^{\langle 2\rangle}(x+1)-\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]^2\Big\}
\end{aligned}$$

from L'Hôspital rule, we get Theorem 12 to complete the proof. □

Taking $x=p$ with $p\in\mathbb{N}_0$ in Theorem 12 and utilizing (9), we deduce the summation formula on harmonic numbers of 2-order.

Corollary 13. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}H_{p+k}^{\langle 2\rangle}=\frac{\binom{p+2n+1}{n}}{2}\Big\{H_{p+2n+1}^{\langle 2\rangle}+H_p^{\langle 2\rangle}-A_n(p)-\tfrac{1}{(p+1)^2}\Big\},$$

where the symbol on the right hand side stands for

$$A_n(p)=\begin{cases}(H_{2q+2n+1}-H_{q+n}-H_{2q+1}+H_q)\\ \times(H_{2q+2n+1}-H_{q+n}-H_{2q+1}+H_q+\frac{2}{2q+1}),\ \ p=2q;\\ (H_{2q+2n+2}-H_{q+n+1}-H_{2q+2}+H_{q+1})\\ \times(H_{2q+2n+2}-H_{q+n+1}-H_{2q+2}+H_q),\qquad p=2q+1.\end{cases}$$

When $p=0$, Corollary 13 reduces to the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}H_k^{\langle 2\rangle}=\frac{\binom{2n+1}{n}}{2}\big\{H_{2n+1}^{\langle 2\rangle}-(H_{2n+1}-H_n)^2\big\}.$$

Lemma 14. *Let a and b be both complex numbers. Then*

$$\begin{aligned}\sum_{k=0}^{n}k\frac{(a)_k(b-a)_k(-n)_k}{k!(\frac{b}{2})_k(-2n)_k}&=\frac{a(b-a)(2a-b+2n)}{b(b+2)}\frac{(\frac{1+a}{2})_n(\frac{2+b-a}{2})_n}{(\frac{1}{2})_n(\frac{4+b}{2})_n}\\&+\frac{a(b-a)(b-2a+2n)}{b(b+2)}\frac{(\frac{2+a}{2})_n(\frac{1+b-a}{2})_n}{(\frac{1}{2})_n(\frac{4+b}{2})_n}.\end{aligned}$$

Proof. For two complex variables x and q, define the q-shifted factorial to be

$$(x;q)_0=1 \quad \text{and} \quad (x;q)_n=\prod_{k=0}^{n-1}(1-xq^k) \quad \text{with} \quad n\in\mathbb{N}.$$

Following Gasper and Rahman [5], define the basic hypergeometric series by

$${}_r\phi_s\left[\begin{matrix}c_1,c_2,\cdots,c_r\\ d_1,d_2,\cdots,d_s\end{matrix}\,\Big|\,q;z\right]=\sum_{k=0}^{\infty}\left[\begin{matrix}c_1,c_2,\cdots,c_r\\ d_1,d_2,\cdots,d_s\end{matrix}\,\Big|\,q\right]_k\left\{(-1)^kq^{\binom{k}{2}}\right\}^{1+s-r}\frac{z^k}{(q;q)_k},$$

where $\{c_i\}_{i\geq 0}$ and $\{d_j\}_{j\geq 1}$ are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Employing the substitutions $a\to q^a$, $b\to q^{1+b}$ in the equation

$${}_4\phi_3\left[\begin{matrix}a,b/a,q^{-n},-q^{-n-1}\\ \sqrt{qb},-\sqrt{qb},q^{-2n-1}\end{matrix}\,\Big|\,q;q\right]=\frac{(qa;q^2)_{n+1}(qb/a;q^2)_{n+1}}{(q;q^2)_{n+1}(qb;q^2)_{n+1}}-q^{n+1}\frac{(a;q^2)_{n+1}(b/a;q^2)_{n+1}}{(q;q^2)_{n+1}(qb;q^2)_{n+1}}$$

due to Wei et al. [17, Corollary 11] and then letting $q\to 1$, we gain

$${}_3F_2\left[\begin{matrix}a,1+b-a,-n\\ \frac{2+b}{2},-2n-1\end{matrix}\,\Big|\,1\right]=\frac{(\frac{1+a}{2})_{n+1}(\frac{2+b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}-\frac{(\frac{a}{2})_{n+1}(\frac{1+b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}.$$

Substitute $1+a$ for a in it to achieve

$${}_3F_2\left[\begin{matrix}1+a,b-a,-n\\ \frac{2+b}{2},-2n-1\end{matrix}\,\Big|\,1\right]=\frac{(\frac{2+a}{2})_{n+1}(\frac{1+b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}-\frac{(\frac{1+a}{2})_{n+1}(\frac{b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}.$$

The linear combination of the last two equations provides

$$\begin{aligned}{}_3F_2\left[\begin{matrix}a,b-a,-n\\ \frac{b}{2},-2n-1\end{matrix}\,\Big|\,1\right]&=\frac{2a-b+2+2n}{b}\frac{(\frac{a}{2})_{n+1}(\frac{1+b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}\\&\quad+\frac{b-2a+2+2n}{b}\frac{(\frac{1+a}{2})_{n+1}(\frac{b-a}{2})_{n+1}}{(\frac{1}{2})_{n+1}(\frac{2+b}{2})_{n+1}}.\end{aligned}\tag{10}$$

It is not difficult to verify that

$$\begin{aligned}\sum_{k=0}^{n}k\frac{(a)_k(b-a)_k(-n)_k}{k!(\frac{b}{2})_k(-2n)_k}&=\sum_{k=1}^{n}\frac{(a)_k(b-a)_k(-n)_k}{(k-1)!(\frac{b}{2})_k(-2n)_k}\\&=\sum_{k=0}^{n-1}\frac{(a)_{k+1}(b-a)_{k+1}(-n)_{k+1}}{k!(\frac{b}{2})_{k+1}(-2n)_{k+1}}\\&=\frac{a(b-a)}{b}{}_3F_2\left[\begin{matrix}1+a,1+b-a,1-n\\ \frac{2+b}{2},1-2n\end{matrix}\,\Big|\,1\right].\end{aligned}$$

Computing the series on the right hand side by (10), we attain Lemma 14 to finish the proof. □

Theorem 15. *Let x be a complex number. Then*

$$\begin{aligned}&\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}kH_k^{\langle 2\rangle}(x)=\frac{x+1}{8}\binom{x+2n+1}{n-1}\\&\times\Big\{4H_{2n+1}^{\langle 2\rangle}(x)-\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})+\tfrac{4}{n}\big]+\tfrac{4}{(x+1)^2}\Big\}.\end{aligned}$$

Proof. The case $a = x+1$ and $b = y+2$ of Lemma 14 can be written as

$$\sum_{k=0}^{n} k\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}} = \frac{4^n(x+1)(y-x+1)(2x-y+2n)}{(y+2)(y+4)}\frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}}{\binom{\frac{y+4}{2}+n}{n}} + \frac{4^n(x+1)(y-x+1)(y-2x+2n)}{(y+2)(y+4)}\frac{\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}}{\binom{\frac{y+4}{2}+n}{n}}. \quad (11)$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (11), we obtain

$$\begin{aligned}&\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}k\big\{H_k(x)-H_k(y-x)\big\}\\ &= \frac{4^n}{(y+2)(y+4)}\frac{\varepsilon_n(x,y)}{\binom{\frac{y+4}{2}+n}{n}} + \frac{4^n(x+1)(y-x+1)}{(y+2)(y+4)}\frac{\zeta_n(x,y)}{\binom{\frac{y+4}{2}+n}{n}},\end{aligned}$$

where the corresponding expressions are

$$\begin{aligned}\varepsilon_n(x,y) &= \{2-6x^2+6xy-(y-2)y-2n(2x-y)\}\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}\\ &\quad -\{2-6x^2+6xy-(y-2)y+2n(2x-y)\}\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n},\\ \zeta_n(x,y) &= \frac{(2x-y+2n)\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n}\big[H_n(\frac{x}{2})-H_n(\frac{y-x+1}{2})\big]}{2}\\ &\quad +\frac{(y-2x+2n)\binom{\frac{x+1}{2}+n}{n}\binom{\frac{y-x}{2}+n}{n}\big[H_n(\frac{x+1}{2})-H_n(\frac{y-x}{2})\big]}{2}.\end{aligned}$$

The last equation can be reformulated as

$$\begin{aligned}&\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}k\sum_{i=1}^{k}\frac{1}{(x+i)(y-x+i)}\\ &= \frac{4^n}{(y+2)(y+4)}\frac{1}{\binom{\frac{y+4}{2}+n}{n}}\frac{\varepsilon_n(x,y)}{y-2x} + \frac{4^n(x+1)(y-x+1)}{(y+2)(y+4)}\frac{1}{\binom{\frac{y+4}{2}+n}{n}}\frac{\zeta_n(x,y)}{y-2x}.\end{aligned}$$

Finding the limit $y \to 2x$ of it and exploiting the two relations

$$\begin{aligned}&\mathrm{Lim}_{y\to 2x}\frac{\varepsilon_n(x,y)}{y-2x}\\ &= \mathrm{Lim}_{y\to 2x}\mathcal{D}_y\varepsilon_n(x,y)\\ &= (x+1)^2\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n}\big[H_n(\tfrac{x+1}{2})-H_n(\tfrac{x}{2})+\tfrac{4n}{(x+1)^2}\big],\\ &\mathrm{Lim}_{y\to 2x}\frac{\zeta_n(x,y)}{y-2x}\\ &= \mathrm{Lim}_{y\to 2x}\mathcal{D}_y\zeta_n(x,y)\\ &= \frac{n}{2}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n}\\ &\quad\times\Big\{4H_{2n}^{\langle 2\rangle}(x+1)-\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})+\tfrac{2}{n}\big]\Big\}\end{aligned}$$

from L'Hôspital rule, we get Theorem 15 to complete the proof. □

Selecting $x = p$ with $p \in \mathbb{N}_0$ in Theorem 15 and using (11), we have the summation formula on harmonic numbers of 2-order.

Corollary 16. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}kH_{p+k}^{\langle 2\rangle}=\frac{p+1}{2}\binom{p+2n+1}{n-1}\Big\{H_{p+2n+1}^{\langle 2\rangle}+H_{p+1}^{\langle 2\rangle}-B_n(p)\Big\},$$

where the symbols on the right hand side stand for

$$B_n(p)=\begin{cases}(H_{2q+2n+1}-H_{q+n}-H_{2q+1}+H_q)\\ \times(H_{2q+2n+1}-H_{q+n}-H_{2q+1}+H_q-\frac{2}{n}), & p=2q;\\ (H_{2q+2n+2}-H_{q+n+1}-H_{2q+2}+H_{q+1})\\ \times(H_{2q+2n+2}-H_{q+n+1}-H_{2q+2}+H_{q+1}+\frac{2}{n}), & p=2q+1.\end{cases}$$

When $p=0$, Corollary 16 produces the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}kH_k^{\langle 2\rangle}=\frac{\binom{2n+1}{n-1}}{2}\big\{H_{2n+1}^{\langle 2\rangle}-\tfrac{2}{n}-(H_{2n+1}-H_n)(H_{2n+1}-H_n-2-\tfrac{2}{n})\big\}.$$

Lemma 17. *Let a and b be both complex numbers. Then*

$$\sum_{k=0}^{n}k^2\frac{(a)_k(b-a)_k(-n)_k}{k!(\frac{b}{2})_k(-2n)_k}=\Theta_n(a,b)\frac{(\frac{1+a}{2})_n(\frac{b-a}{2})_{n+1}}{(\frac{1}{2})_n(\frac{4+b}{2})_{n+1}}+\Omega_n(a,b)\frac{(\frac{a}{2})_{n+1}(\frac{1+b-a}{2})_n}{(\frac{1}{2})_n(\frac{4+b}{2})_{n+1}}.$$

where the corresponding expressions are

$$\Theta_n(a,b)=\frac{2a^3(2a-3b+2n)+2a^2(b^2-b-3bn-2n^2+2n+2)+a(2n+1)(b^2-2b+2bn+4n)}{b(b+2)},$$
$$\Omega_n(a,b)=\frac{(a-b)^3(2a+1)+2(a-b)^2(a^2-an+n+1)-(a-b)(2n+1)(a^2-2a+2an+4n)}{b(b+2)}.$$

Proof. Performing the replacements $a\to q^a$, $b\to q^{1+b}$ in the equation

$${}_4\phi_3\left[\begin{matrix}a,b/a,q^{-n},-q^{-n-2}\\ \sqrt{qb},-\sqrt{qb},q^{-2n-2}\end{matrix}\,\Big|\,q;q\right]=\frac{(1+q)(1-q^{2n+3})}{q^{n+1}-q^{-n-1}}\frac{(a;q^2)_{n+2}(b/a;q^2)_{n+2}}{(q;q^2)_{n+2}(qb;q^2)_{n+2}}$$
$$+\left\{1-\frac{(1-a)(1-b/a)(1-q^{2n+4})}{(1-q^{-2n-2})(1-aq^{2n+3})(1-bq^{2n+3}/a)}\right\}\frac{(qa;q^2)_{n+2}(qb/a;q^2)_{n+2}}{(q;q^2)_{n+2}(qb;q^2)_{n+2}}.$$

due to Wei et al. [17, Corollary 12] and then letting $q\to 1$, we gain

$${}_3F_2\left[\begin{matrix}a,1+b-a,-n\\ \frac{2+b}{2},-2n-2\end{matrix}\,\Big|\,1\right]=-\frac{2n+3}{n+1}\frac{(\frac{a}{2})_{n+2}(\frac{1+b-a}{2})_{n+2}}{(\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}$$
$$+\frac{a(1+b-a)(n+2)+(n+1)(a+2n+3)(b-a+2n+4)}{(n+1)(a+2n+3)(b-a+2n+4)}\frac{(\frac{1+a}{2})_{n+2}(\frac{2+b-a}{2})_{n+2}}{(\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}.$$

Replace a by $1+a$ in it to achieve

$${}_3F_2\left[\begin{matrix}1+a,b-a,-n\\ \frac{2+b}{2},-2n-2\end{matrix}\,\Big|\,1\right]=-\frac{2n+3}{n+1}\frac{(\frac{1+a}{2})_{n+2}(\frac{b-a}{2})_{n+2}}{(\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}$$
$$+\frac{(1+a)(b-a)(n+2)+(n+1)(a+2n+4)(b-a+2n+3)}{(n+1)(a+2n+4)(b-a+2n+3)}\frac{(\frac{2+a}{2})_{n+2}(\frac{1+b-a}{2})_{n+2}}{(\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}.$$

The linear combination of the last two equations offers

$${}_3F_2\left[\begin{matrix}a,b-a,-n\\ \frac{b}{2},-2n-2\end{matrix}\,\Big|\,1\right]=\frac{a(2a-b+n+1)-(n+1)(b-a+2n+4)}{(n+1)(a-1)b}\frac{(\frac{a-1}{2})_{n+2}(\frac{b-a}{2})_{n+2}}{(-\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}$$
$$+\frac{(b-a)(b-2a+n+1)-(n+1)(a+2n+4)}{(n+1)(b-a-1)b}\frac{(\frac{a}{2})_{n+2}(\frac{b-a-1}{2})_{n+2}}{(-\frac{1}{2})_{n+2}(\frac{2+b}{2})_{n+2}}.\qquad(12)$$

It is routine to show that

$$\begin{aligned}
&\sum_{k=0}^{n} k^2 \frac{(a)_k (b-a)_k (-n)_k}{k!(\frac{b}{2})_k (-2n)_k} = \sum_{k=1}^{n} k \frac{(a)_k (b-a)_k (-n)_k}{(k-1)!(\frac{b}{2})_k (-2n)_k} \\
&= \sum_{k=0}^{n-1} (k+1) \frac{(a)_{k+1} (b-a)_{k+1} (-n)_{k+1}}{k!(\frac{b}{2})_{k+1} (-2n)_{k+1}} \\
&= \frac{a(b-a)}{b} {}_3F_2\left[\begin{matrix} 1+a, 1+b-a, 1-n \\ \frac{2+b}{2}, 1-2n \end{matrix} \,\middle|\, 1\right] \\
&+ \frac{a(b-a)}{b} \sum_{k=1}^{n-1} \frac{(1+a)_k (1+b-a)_k (1-n)_k}{(k-1)!(\frac{2+b}{2})_k (1-2n)_k} \\
&= \frac{a(b-a)}{b} {}_3F_2\left[\begin{matrix} 1+a, 1+b-a, 1-n \\ \frac{2+b}{2}, 1-2n \end{matrix} \,\middle|\, 1\right] \\
&+ \frac{2a(1+a)(b-a)(1+b-a)(1-n)}{b(2+b)(1-2n)} {}_3F_2\left[\begin{matrix} 2+a, 2+b-a, 2-n \\ \frac{4+b}{2}, 2-2n \end{matrix} \,\middle|\, 1\right].
\end{aligned}$$

Evaluating the two series on the right hand side by (10) and (12), we attain Lemma 17 to finish the proof. □

Theorem 18. *Let x be a complex number. Then*

$$\begin{aligned}
&\sum_{k=0}^{n} \binom{2n-k}{n} \binom{x+k}{k} k^2 H_k^{\langle 2\rangle}(x) = \frac{(x+1)(xn+3n+1)}{8(n-1)} \binom{x+2n+1}{n-2} \\
&\times \Big\{ 4H_{2n+1}^{\langle 2\rangle}(x) + \tfrac{8(x^2+3x+n^2-n+1)}{n(x+1)(xn+3n+1)} - \big[H_n(\tfrac{x}{2}) - H_{n+1}(\tfrac{x-1}{2})\big] \\
&\quad \times \big[H_n(\tfrac{x}{2}) - H_{n+1}(\tfrac{x-1}{2}) - \tfrac{4(x+n+1)(x^2+3x-2n+1)}{n(x+1)(xn+3n+1)}\big] \Big\}.
\end{aligned}$$

Proof. The case $a = x+1$ and $b = y+2$ of Lemma 17 can be manipulated as

$$\sum_{k=0}^{n} k^2 \frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}} = \delta_n(x,y) \frac{\binom{\frac{x+1}{2}+n}{n+1}\binom{\frac{y-x}{2}+n}{n}}{\binom{\frac{y+6}{2}+n}{n+1}} + \xi_n(x,y) \frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n+1}}{\binom{\frac{y+6}{2}+n}{n+1}}, \quad (13)$$

where the symbols on the right hand side stand for

$$\begin{aligned}
\delta_n(x,y) &= \frac{4^n(x-y-1)}{y+2} \frac{2x^2(2x-3y-2n)+2x(y^2-3y+yn-2-2n^2)+3y^2+2y-4n-12n^2}{y+4}, \\
\xi_n(x,y) &= \frac{4^n(x+1)}{y+2} \frac{2x^2(2x-3y+2n)+2x(y^2-3y-3yn-2-2n^2)+(2n+3)y^2+2(2n^2+1)y+4n+12n^2}{y+4}.
\end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of it, we obtain

$$\sum_{k=0}^{n} \frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}} k^2 \big\{H_k(x) - H_k(y-x)\big\} = \frac{\varphi_n(x,y) + \psi_n(x,y)}{\binom{\frac{y+6}{2}+n}{n+1}},$$

where the the corresponding expressions are

$$\begin{aligned}
\varphi_n(x,y) &= \mathcal{D}_x[\delta_n(x,y)] \binom{\frac{x+1}{2}+n}{n+1} \binom{\frac{y-x}{2}+n}{n} + D_x[\xi_n(x,y)] \binom{\frac{x}{2}+n}{n} \binom{\frac{y-x+1}{2}+n}{n+1}, \\
\psi_n(x,y) &= \frac{\delta_n(x,y)\binom{\frac{x+1}{2}+n}{n+1}\binom{\frac{y-x}{2}+n}{n}\big[H_{n+1}(\frac{x-1}{2}) - H_n(\frac{y-x}{2})\big]}{2} \\
&\quad + \frac{\xi_n(x,y)\binom{\frac{x}{2}+n}{n}\binom{\frac{y-x+1}{2}+n}{n+1}\big[H_n(\frac{x}{2}) - H_{n+1}(\frac{y-x-1}{2})\big]}{2}.
\end{aligned}$$

The last equation can be reformulated as

$$\sum_{k=0}^{n}\frac{\binom{x+k}{k}\binom{y-x+k}{k}\binom{2n-k}{n}}{\binom{\frac{y}{2}+k}{k}}k^2\sum_{i=1}^{k}\frac{1}{(x+i)(y-x+i)}$$
$$=\frac{1}{\binom{\frac{y+6}{2}+n}{n+1}}\frac{\varphi_n(x,y)}{y-2x}+\frac{1}{\binom{\frac{y+6}{2}+n}{n+1}}\frac{\psi_n(x,y)}{y-2x}.$$

Fining the limit $y\to 2x$ of it and utilizing the two relations

$$\begin{aligned}
&\mathrm{Lim}_{y\to 2x}\frac{\varphi_n(x,y)}{y-2x}\\
&=\mathrm{Lim}_{y\to 2x}\mathcal{D}_y\varphi_n(x,y)\\
&=\frac{4^n(x+n+1)(x^2+3x-2n+1)}{2(x+1)(x+2)}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n+1}\\
&\times\ \big[H_n(\tfrac{x}{2})-H_{n+1}(\tfrac{x-1}{2})+\tfrac{4(x^2+3x+n^2-n+1)}{(x+n+1)(x^2+3x-2n+1)}\big],\\
&\mathrm{Lim}_{y\to 2x}\frac{\psi_n(x,y)}{y-2x}\\
&=\mathrm{Lim}_{y\to 2x}\mathcal{D}_y\psi_n(x,y)\\
&=\frac{4^{n-1}n(xn+3n+1)}{x+2}\binom{\frac{x}{2}+n}{n}\binom{\frac{x+1}{2}+n}{n+1}\Big\{4H^{\langle 2\rangle}_{2n+1}(x)\\
&-\ \big[H_n(\tfrac{x}{2})-H_{n+1}(\tfrac{x-1}{2})\big]\big[H_n(\tfrac{x}{2})-H_{n+1}(\tfrac{x-1}{2})-\tfrac{2(x+n+1)(x^2+3x-2n+1)}{n(x+1)(xn+3n+1)}\big]\Big\}
\end{aligned}$$

from L'Hôspital rule, we get Theorem 18 to complete the proof. □

Fixing $x=p$ with $p\in\mathbb{N}_0$ in Theorem 18 and exploiting (13), we deduce the summation formula on harmonic numbers of 2-order.

Corollary 19. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}k^2H^{\langle 2\rangle}_{p+k}=\frac{(p+1)(pn+3n+1)}{2(n-1)}\binom{p+2n+1}{n-2}$$
$$\times\Big\{H^{\langle 2\rangle}_{p+2n+1}+H^{\langle 2\rangle}_{p}-C_n(p)+\tfrac{2(p^2+3p+n^2-n+1)}{n(p+1)(pn+3n+1)}\Big\},$$

where the symbol on the right hand side stands for

$$C_n(p)=\begin{cases}
\big[H_{2q+2n+1}-H_{q+n}-H_{2q}+H_q\big]\\
\times\big[H_{2q+2n+1}-H_{q+n}-H_{2q}+H_q+\frac{2(2q+n+1)(4q^2+6q-2n+1)}{n(2q+1)(2qn+3n+1)}\big], & p=2q;\\
\big[H_{2q+2n+2}-H_{q+n+1}-H_{2q+1}+H_q\big]\\
\times\big[H_{2q+2n+2}-H_{q+n+1}-H_{2q+1}+H_q-\frac{(2q+n+2)(4q^2+10q-2n+5)}{n(q+1)(2qn+4n+1)}\big], & p=2q+1.
\end{cases}$$

When $p=0$, Corollary 19 creates the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}k^2H^{\langle 2\rangle}_{k}=\frac{3n+1}{2(n-1)}\binom{2n+1}{n-2}$$
$$\times\Big\{H^{\langle 2\rangle}_{2n+1}-\big[H_{2n+1}-H_n\big]\big[H_{2n+1}-H_n-\tfrac{2(n+1)(2n-1)}{n(3n+1)}\big]+\tfrac{2(n^2-n+1)}{n(3n+1)}\Big\}.$$

4. The third family of summation formulae involving

GENERALIZED HARMONIC NUMBERS

Theorem 20. *Let x be a complex number. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}H_k^2(x)=\frac{\binom{x+2n+1}{n}}{8}\Big\{4H_n^{\langle 2\rangle}(x+1)-4H_n^{\langle 2\rangle}(x+n+1)$$
$$+8H_n^2(x+n+1)-\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})-\tfrac{4}{x+1}\big]\Big\}.$$

Proof. The case $y=2x$ of (9) reads as

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}=\binom{x+2n+1}{n}. \tag{14}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (14), we gain

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}H_k(x)=\binom{x+2n+1}{n}\big\{H_{2n+1}(x)-H_{n+1}(x)\big\}. \tag{15}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (15), we achieve

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}\big\{H_k^2(x)-H_k^{\langle 2\rangle}(x)\big\}$$
$$=\binom{x+2n+1}{n}\big\{[H_{2n+1}(x)-H_{n+1}(x)]^2-[H_{2n+1}^{\langle 2\rangle}(x)-H_{n+1}^{\langle 2\rangle}(x)]\big\}.$$

The sum of Theorem 12 and the last equation gives Theorem 20. □

Selecting $x=p$ with $p\in\mathbb{N}_0$ in Theorem 20 and using (14) and (15), we have the summation formula on harmonic numbers.

Corollary 21. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}H_{p+k}^2=\frac{\binom{p+2n+1}{n}}{2}\Big\{2H_{p+n+1}^{\langle 2\rangle}-H_{p+2n+1}^{\langle 2\rangle}-H_{p+1}^{\langle 2\rangle}$$
$$+2(H_{p+n+1}-H_{p+2n+1})^2-2H_p(2H_{p+n+1}-2H_{p+2n+1}-H_p)-A_n(p)\Big\},$$

where $A_n(p)$ has appeared in Corollary 13.

When $p=0$, Corollary 21 reduces to the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}H_k^2=\frac{\binom{2n+1}{n}}{2}\big\{2H_{n+1}^{\langle 2\rangle}-H_{2n+1}^{\langle 2\rangle}+2H_{n+1}^2-H_n^2$$
$$+H_{2n+1}(H_{2n+1}-4H_{n+1}+2H_n)\big\}.$$

Theorem 22. *Let x be a complex number. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}kH_k^2(x)=\frac{x+1}{8}\binom{x+2n+1}{n-1}\Big\{8H_{n+2}^{\langle 2\rangle}(x)-4H_{2n}^{\langle 2\rangle}(x+1)$$
$$+8\big[H_{n+2}(x)-H_{2n+1}(x)\big]\big[H_{n+2}(x)-H_{2n+1}(x)-\tfrac{2}{x+1}\big]$$
$$-\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})\big]\big[H_n(\tfrac{x}{2})-H_n(\tfrac{x+1}{2})+\tfrac{4}{n}\big]\Big\}.$$

Proof. The case $y=2x$ of (11) can be written as

$$\sum_{k=0}^{n} k\binom{2n-k}{n}\binom{x+k}{k}=(x+1)\binom{x+2n+1}{n-1}. \tag{16}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (16), we attain

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}kH_k(x)=(x+1)\binom{x+2n+1}{n-1}\big\{H_{2n+1}(x)-H_{n+1}(x+1)\big\}. \tag{17}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (17), we obtain

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}k\big\{H_k^2(x)-H_k^{\langle 2\rangle}(x)\big\}=(x+1)\binom{x+2n+1}{n-1}$$
$$\times\Big\{[H_{n+2}^{\langle 2\rangle}(x)-H_{2n+1}^{\langle 2\rangle}(x)]+[H_{n+2}(x)-H_{2n+1}(x)][H_{n+2}(x)-H_{2n+1}(x)-\tfrac{2}{x+1}]\Big\}.$$

The sum of Theorem 15 and the last equation provides Theorem 22. □

Choosing $x=p$ with $p\in\mathbb{N}_0$ in Theorem 22 and utilizing (16) and (17), we deduce the summation formula on harmonic numbers.

Corollary 23. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}kH_{p+k}^2=\frac{p+1}{2}\binom{p+2n+1}{n-1}\Big\{2H_{p+n+2}^{\langle 2\rangle}-H_{p+2n+1}^{\langle 2\rangle}-H_{p+1}^{\langle 2\rangle}$$
$$+2(H_{p+n+2}-H_{p+2n+1})^2-2H_{p+1}(2H_{p+n+2}-2H_{p+2n+1}-H_{p+1})-B_n(p)\Big\},$$

where $B_n(p)$ has emerged in Corollary 16.

When $p=0$, Corollary 23 produces the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}kH_k^2=\frac{\binom{2n+1}{n-1}}{2}\Big\{2H_{n+2}^{\langle 2\rangle}-H_{2n+1}^{\langle 2\rangle}+\big[H_{2n+1}-H_{n+2}\big]$$
$$\times\big[H_{2n+1}-H_{n+2}+\tfrac{2(3n^3+8n^2+6n+2)}{n(n+1)(n+2)}\big]+\tfrac{2n^4+6n^3+6n^2+5n+4}{n(n+1)^2(n+2)^2}\Big\}.$$

Theorem 24. *Let x be a complex number. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}k^2H_k^2(x)=\frac{(x+1)(xn+3n+1)}{8(n-1)}\binom{x+2n+1}{n-2}$$
$$\times\Big\{8\big[H_{n+3}(x)-H_{2n+1}(x)\big]\big[H_{n+3}(x)-H_{2n+1}(x)-\tfrac{2(2xn+4n+1)}{(x+1)(xn+3n+1)}\big]$$
$$-\big[H_n(\tfrac{x}{2})-H_{n+1}(\tfrac{x-1}{2})\big]\big[H_n(\tfrac{x}{2})-H_{n+1}(\tfrac{x-1}{2})-\tfrac{4(x+n+1)(x^2+3x-2n+1)}{n(x+1)(xn+3n+1)}\big]$$
$$+8H_{n+3}^{\langle 2\rangle}(x)-4H_{2n+1}^{\langle 2\rangle}(x)+\tfrac{8(x^2+3x+3n^2-n+1)}{n(x+1)(xn+3n+1)}\Big\}.$$

Proof. The case $y=2x$ of (13) can be manipulated as

$$\sum_{k=0}^{n}k^2\binom{2n-k}{n}\binom{x+k}{k}=\frac{(x+1)(xn+3n+1)}{n-1}\binom{x+2n+1}{n-2}. \tag{18}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (18), we get

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}k^2H_k(x)=\frac{(x+1)(xn+3n+1)}{n-1}$$
$$\times\binom{x+2n+1}{n-2}\Big\{H_{2n+1}(x)-H_{n+2}(x+1)+\tfrac{n}{xn+3n+1}\Big\}. \tag{19}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (19), we gain

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{x+k}{k}k^2\big\{H_k^2(x)-H_k^{\langle 2\rangle}(x)\big\}=\frac{(x+1)(xn+3n+1)}{n-1}$$
$$\times\binom{x+2n+1}{n-2}\Big\{H_{n+3}^{\langle 2\rangle}(x)-H_{2n+1}^{\langle 2\rangle}(x)+\big[H_{n+3}(x)-H_{2n+1}(x)\big]$$
$$\times\big[H_{n+3}(x)-H_{2n+1}(x)-\tfrac{2(2xn+4n+1)}{(x+1)(xn+3n+1)}\big]+\tfrac{2n}{(x+1)(xn+3n+1)}\Big\}.$$

The sum of Theorem 15 and the last equation offers Theorem 24. □

Taking $x=p$ with $p\in\mathbb{N}_0$ in Theorem 24 and exploiting (18) and (19), we have the summation formula on harmonic numbers.

Corollary 25. *Let p be a nonnegative integer. Then*

$$\sum_{k=0}^{n}\binom{2n-k}{n}\binom{p+k}{k}k^2H_{p+k}^2=\frac{(p+1)(pn+3n+1)}{2(n-1)}\binom{p+2n+1}{n-2}$$
$$\times\Big\{2(H_{p+n+3}-H_{p+2n+1})\big(H_{p+n+3}-H_{p+2n+1}-2H_{p+1}-\tfrac{2n}{pn+3n+1}\big)+2H_p^2$$
$$+\tfrac{4(2pn+4n+1)}{(p+1)(pn+3n+1)}H_p+\tfrac{2(p^2+3p+3n^2-n+1)}{n(p+1)(pn+3n+1)}+2H_{p+n+3}^{\langle 2\rangle}-H_{p+2n+1}^{\langle 2\rangle}-H_p^{\langle 2\rangle}-C_n(p)\Big\},$$

where $C_n(p)$ has been displayed in Corollary 19.

When $p=0$, Corollary 25 creates the concise harmonic number identity:

$$\sum_{k=0}^{n}\binom{2n-k}{n}k^2H_k^2=\frac{3n+1}{2(n-1)}\binom{2n+1}{n-2}\Big\{2\big[H_{2n+1}-H_{n+3}\big]\big[H_{2n+1}-H_{n+3}+\tfrac{8n+2}{3n+1}\big]$$
$$-\big[H_{2n+1}-H_n\big]\big[H_{2n+1}-H_n-\tfrac{2(n+1)(2n-1)}{n(3n+1)}\big]+2H_{n+3}^{\langle 2\rangle}-H_{2n+1}^{\langle 2\rangle}+\tfrac{2(3n^2-n+1)}{n(3n+1)}\Big\}.$$

Remark: (15), (17) and (19) can also be derived by means of the derivative operator and Chu-Vandermonde convolution:

$$\sum_{k=0}^{n}\binom{x}{k}\binom{y}{n-k}=\binom{x+y}{n}.$$

The reader may refer to Wei et. al. [18] for details. Some similar results can be seen in Wei et. al. [19].

Acknowledgments

The work is supported by the National Natural Science Foundation of China (No. 11301120).

References

[1] W.N. Bailey, On the sum of a terminating ${}_3F_2$, Quart. J. Math. Oxford (2). 4 (1953), 237–240.
[2] G.E. Andrews, K. Uchimura, Identities in combinatorics IV: differentiation and harmonic numbers, Utilitas Math. 28 (1985), 265–269.
[3] G.E. Andrews, R. Askey, R. Roy, Special Functions, Cambridge University Press, Cambridge, 2000.
[4] Y. Chen, Q. Hou, H. Jin, The Abel-Zeilberger algorithm, Electron. J. Comb. 18 (2011), #P17.
[5] G. Gasper, M. Rahman, Basic Hypergeometric Series (2nd edition), Cambridge Univercity Press, Cambridge, 2004.
[6] C. Krattenthaler, T. Rivoal, Hypergéométrie et fonction zêta de Riemann, Mem. Amer. Math. Soc. 186, no. 875, Providence, R. I., 2007.

[7] M.J. Kronenburg, Some generalized harmonic number identities, arXiv:1103.5430v2 [math.NT], 2012.
[8] M.J. Kronenburg, On two types of harmonic number identities, arXiv:1202.3981v2 [math.NT], 2012.
[9] P. Paule, C. Schneider, Computer proofs of a new family of harmonic number identities, Adv. Appl. Math. 31 (2003), 359–378.
[10] C. Schneider, Symbolic summation assists Combinatorics, Sém. Lothar. Combin. 56 (2006), Article B56b.
[11] A. Sofo, Sums of derivatives of binomial coefficients, Adv. Appl. Math. 42 (2009), 123–134.
[12] A. Sofo, Quadratic alternating harmonic number sums, J. Number Theory 154 (2015), 144–159.
[13] Z. Sun, Arithmetic theory of harmonic numbers, Proc. Amer. Math. Soc. 140 (2012), 415–428.
[14] W. Wang, Riordan arrays and harmonic number identities, Comput. Math. Appl. 60 (2010), 1494–1509.
[15] W. Wang, C. Jia, Harmonic number identities via the Newton-Andrews method, Ramanujan J. 35 (2014), 263–285.
[16] J. Wang, C. Wei, Derivative operator and summation formulae involving generalized harmonic numbers, J. Math. Anal. Appl. 434 (2016) 315–341.
[17] C. Wei, D. Gong, J. Li, Summation formulae for q-Watson type ${}_4\phi_3$-series, Discrete Math. 313 (2013), 1589–1593.
[18] C. Wei, D. Gong, Q. Wang, Chu-Vandermonde convolution and harmonic number identities, Integral Transforms Spec. Funct. 24 (2013), 324–330.
[19] C. Wei, D. Gong, Q. Yan, Telescoping method, derivative operators and harmonic number identities, Integral Transforms Spec. Funct. 25 (2014), 203–214.